



FINAL REPORT

Part 1 - Summary Details

Cotton CRC Project Number: 1.04.11.04

Project Title: BMP Border Rivers

Project Commencement Date: 1.6.2008 **Project Completion Date:** 31.5.2012

Cotton CRC Program: 1.4 Resilient Farming

Part 2 – Contact Details

Administrator: Brendan Griffiths

Organisation: Griffiths Agriculture

Postal Address:	P.O. Box 1044 Goondiwindi, 4390
------------------------	---------------------------------

Ph: 07 46715990

Fax: 07 46715991

E-mail: griffb@bigpond.com

Principal Researcher: Brendan Griffiths

Organisation: Griffiths Agriculture

Postal Address:	P.O. Box 1044 Goondiwindi
------------------------	---------------------------

Ph: 07 46715990

Fax: 07 46715991

E-mail: griffb@bigpond.com

Supervisor: Graham Harris

Organisation: DEEDI, Qld

Postal Address:	
------------------------	--

Ph: 07 4688 1559

Fax: 07 4688 1199

E-mail: Graham.Harris@deedi.qld.gov.au

Signature of Research Provider Representative: _____

Part 3 – Final Report Guide (due at 31st May 2012)

Background

BMP Border Rivers was a collaboration between the projects High Yielding Wheat in Cotton Farming Systems, and PhD Brendan Griffiths, and the private consultancy Griffiths Agriculture to extend research pertaining to the development of a system of achieving high yielding wheat in cotton farming systems in the Border Rivers region of Northern NSW, and Southern Queensland.

Objectives

To investigate / compare our current farming systems with respect to southern management systems for irrigated cereals, with specific reference to the northern farming systems associated with a cotton / cereal rotation.

Many questions have been raised regarding management and farming systems associated with intensive rotation with cotton and cereal rotations, with respect to such issues as;

- Canopy management.
- Appropriate row configuration.
- Beds vs hills.
- Is zero/ minimum till possible, and what benefit is there with stubble retention?
- How can we practically get into cereal cotton rotations and fulfil such obligations as pupal destruction?

Methods

The methods utilised throughout the course of this project have been clearly outlined in the reporting process in Centric. Working closely with PhD Brendan Griffiths and High Yielding Grains in Cotton Farming systems, scientifically rigorous experiments were designed to both investigate and demonstrate methods being employed in the development of a system achieving high yielding wheat, in a cotton system. The other components of the extension effort included the development of a field demonstration site, planted in each year of the project to demonstrate the applied methods of the research being generated through the High Yielding Grains in Cotton Farming Systems Project. Two field walks were then conducted at this site attracting key industry personnel interested in the development of a high yielding wheat cropping system specific to the northern cropping region of NSW, and Southern Queensland.

Results

Please see attached document "Irrigated Wheat - Best Practice Guidelines in Cotton Farming Systems".

Conclusion

There have been several key take home messages that have been found by this project. The first would be that the Southern systems that have previously been developed for the production of high yielding irrigated wheat are not relevant to those in the northern region of NSW and Southern Queensland. This was largely as a result of the obvious climatic differences, principally the milder start to the growing season, and a shorter end to the season. There are several other take home messages clearly outlined in the attached publication "Irrigated Wheat - Best Practice Guidelines in Cotton Farming Systems".

Extension Opportunities

As a result of this project there will be many opportunities to further the extension work delivered throughout the course of this project. See the attached publication "Irrigated wheat - Best practice guidelines in Cotton Farming Systems". There has been an entire cropping program devised throughout the course of this project pertinent to the achievement of high yielding wheats in the northern cropping environment of NSW, and that of Southern Queensland.

Publications

Please refer to uploads on Centric for following documents.

Presentations (conference, field days, workshops etc)

Wheat Systems 2010 N Water Trial.doc
PresentationGriffithsBrendanwheatGNDI.ppt
Wheat Systems 2010 Row Spacing Trial.doc
Cotton CRC review 2010 BBG.ppt
HYGICFS Project Meeting Sept 2010.ppt
Table 2 ModdusInWheatInterimData.xls
table 1 WheatSystemsYieldMeans2011.xls

Non-peered reviewed articles

review of mechanisms BBG301209#2.doc
Irrigated Wheat Literature Review BBG020609.doc
Draft Agnote BBG240310.doc
Enhanced P Protocol 20101.doc
abstract crc reviewBBG151010.doc
HYIGCFS 2011 PLAN Var RequestNOV 2010.doc
Protocol 2 WP032B3-2011AU - TXP WHEAT ROOT ENHANCEMENT FIELD.doc
Protocol 4 WheatSystems2011NTrial.doc
Protocol 1 WP032D3-2011AU - MODDUS WHEAT WSC FIELD.doc
Protocol 5 WP032E3-2011AU - MODDUS WHEAT WSC+ROOT EN GHOUSE__.doc
Protocol 3 WheatSystems2011RowSpacingTrial.doc

Part 4 – Final Report Executive Summary

BMP Border Rivers was a collaboration between the projects High Yielding Wheat in Cotton Farming Systems, and PhD Brendan Griffiths, and the private consultancy Griffiths Agriculture to extend research pertaining to the development of a system of achieving high yielding wheat in cotton farming systems in the Border Rivers region of Northern NSW, and Southern Queensland.

Originally designed to implement practices from Southern irrigated farming systems, the project team very early in the project realised that these practices would not be directly applicable to a northern farming system, principally as a result of the interaction between crop physiology, and the obvious climatic differences, being a milder start to the season, and a hotter, shorter finish to the growing season. As such the project team set about developing a system that would be directly applicable to the northern irrigated cropping system, and environment, with very good success.

Yields achieved at the start of the project, using methods developed in the southern high yielding grains projects, were rarely in excess of 6 t/ha. There were also associated issues with respects to crop physiology and grain quality. By the end of the project, experimental yields in excess of 9t/ha were able to be achieved, using improved methods of managing crop canopy, and other agronomic methods to manipulate various components of crop physiology, with the view of maximising yields, and minimising lodging and the associated grain quality problems.

Throughout the course of the project a field demonstration site, displaying the research and methods discovered in the High Yielding Grains in Cotton Farming Systems and in PhD Brendan Griffiths projects, was set up in each year of the project. There were two field walks conducted at this site in each year of the project attracting relevant interested industry personnel interested in the development of best practice guidelines for the consistent achievement of high yielding wheats in a cotton farming system.

A publication titled “Irrigated Wheat- Best Practice Guidelines in Cotton Farming Systems” was generated as part of the project outlining a system that will more

consistently allow producers in the northern growing region of NSW, and Southern Queensland, the ability to achieve high yielding, high quality wheats in an irrigated cotton farming system.



FINAL REPORT

Part 1 - Summary Details

Cotton CRC Project Number: 1.04.11.04

Project Title: BMP Border Rivers

Project Commencement Date: 1.6.2008 **Project Completion Date:** 31.5.2012

Cotton CRC Program: 1.4 Resilient Farming

Part 2 – Contact Details

Administrator: Brendan Griffiths

Organisation: Griffiths Agriculture

Postal Address:	P.O. Box 1044 Goondiwindi, 4390
------------------------	---------------------------------

Ph: 07 46715990

Fax: 07 46715991

E-mail: griffb@bigpond.com

Principal Researcher: Brendan Griffiths

Organisation: Griffiths Agriculture

Postal Address:	P.O. Box 1044 Goondiwindi
------------------------	---------------------------

Ph: 07 46715990

Fax: 07 46715991

E-mail: griffb@bigpond.com

Supervisor: Graham Harris

Organisation: DEEDI, Qld

Postal Address:	
------------------------	--

Ph: 07 4688 1559

Fax: 07 4688 1199

E-mail: Graham.Harris@deedi.qld.gov.au

Signature of Research Provider Representative: _____

Part 3 – Final Report Guide (due at 31st May 2012)

Background

BMP Border Rivers was a collaboration between the projects High Yielding Wheat in Cotton Farming Systems, and PhD Brendan Griffiths, and the private consultancy Griffiths Agriculture to extend research pertaining to the development of a system of achieving high yielding wheat in cotton farming systems in the Border Rivers region of Northern NSW, and Southern Queensland.

Objectives

To investigate / compare our current farming systems with respect to southern management systems for irrigated cereals, with specific reference to the northern farming systems associated with a cotton / cereal rotation.

Many questions have been raised regarding management and farming systems associated with intensive rotation with cotton and cereal rotations, with respect to such issues as;

- Canopy management.
- Appropriate row configuration.
- Beds vs hills.
- Is zero/ minimum till possible, and what benefit is there with stubble retention?
- How can we practically get into cereal cotton rotations and fulfil such obligations as pupal destruction?

Methods

The methods utilised throughout the course of this project have been clearly outlined in the reporting process in Centric. Working closely with PhD Brendan Griffiths and High Yielding Grains in Cotton Farming systems, scientifically rigorous experiments were designed to both investigate and demonstrate methods being employed in the development of a system achieving high yielding wheat, in a cotton system. The other components of the extension effort included the development of a field demonstration site, planted in each year of the project to demonstrate the applied methods of the research being generated through the High Yielding Grains in Cotton Farming Systems Project. Two field walks were then conducted at this site attracting key industry personnel interested in the development of a high yielding wheat cropping system specific to the northern cropping region of NSW, and Southern Queensland.

Results

Please see attached document "Irrigated Wheat - Best Practice Guidelines in Cotton Farming Systems".

Conclusion

There have been several key take home messages that have been found by this project. The first would be that the Southern systems that have previously been developed for the production of high yielding irrigated wheat are not relevant to those in the northern region of NSW and Southern Queensland. This was largely as a result of the obvious climatic differences, principally the milder start to the growing season, and a shorter end to the season. There are several other take home messages clearly outlined in the attached publication "Irrigated Wheat - Best Practice Guidelines in Cotton Farming Systems".

Extension Opportunities

As a result of this project there will be many opportunities to further the extension work delivered throughout the course of this project. See the attached publication "Irrigated wheat - Best practice guidelines in Cotton Farming Systems". There has been an entire cropping program devised throughout the course of this project pertinent to the achievement of high yielding wheats in the northern cropping environment of NSW, and that of Southern Queensland.

Publications

Please refer to uploads on Centric for following documents.

Presentations (conference, field days, workshops etc)

Wheat Systems 2010 N Water Trial.doc
PresentationGriffithsBrendanwheatGNDI.ppt
Wheat Systems 2010 Row Spacing Trial.doc
Cotton CRC review 2010 BBG.ppt
HYGICFS Project Meeting Sept 2010.ppt
Table 2 ModdusInWheatInterimData.xls
table 1 WheatSystemsYieldMeans2011.xls

Non-peered reviewed articles

review of mechanisms BBG301209#2.doc
Irrigated Wheat Literature Review BBG020609.doc
Draft Agnote BBG240310.doc
Enhanced P Protocol 20101.doc
abstract crc reviewBBG151010.doc
HYIGCFS 2011 PLAN Var RequestNOV 2010.doc
Protocol 2 WP032B3-2011AU - TXP WHEAT ROOT ENHANCEMENT FIELD.doc
Protocol 4 WheatSystems2011NTrial.doc
Protocol 1 WP032D3-2011AU - MODDUS WHEAT WSC FIELD.doc
Protocol 5 WP032E3-2011AU - MODDUS WHEAT WSC+ROOT EN GHOUSE__.doc
Protocol 3 WheatSystems2011RowSpacingTrial.doc

Part 4 – Final Report Executive Summary

BMP Border Rivers was a collaboration between the projects High Yielding Wheat in Cotton Farming Systems, and PhD Brendan Griffiths, and the private consultancy Griffiths Agriculture to extend research pertaining to the development of a system of achieving high yielding wheat in cotton farming systems in the Border Rivers region of Northern NSW, and Southern Queensland.

Originally designed to implement practices from Southern irrigated farming systems, the project team very early in the project realised that these practices would not be directly applicable to a northern farming system, principally as a result of the interaction between crop physiology, and the obvious climatic differences, being a milder start to the season, and a hotter, shorter finish to the growing season. As such the project team set about developing a system that would be directly applicable to the northern irrigated cropping system, and environment, with very good success.

Yields achieved at the start of the project, using methods developed in the southern high yielding grains projects, were rarely in excess of 6 t/ha. There were also associated issues with respects to crop physiology and grain quality. By the end of the project, experimental yields in excess of 9t/ha were able to be achieved, using improved methods of managing crop canopy, and other agronomic methods to manipulate various components of crop physiology, with the view of maximising yields, and minimising lodging and the associated grain quality problems.

Throughout the course of the project a field demonstration site, displaying the research and methods discovered in the High Yielding Grains in Cotton Farming Systems and in PhD Brendan Griffiths projects, was set up in each year of the project. There were two field walks conducted at this site in each year of the project attracting relevant interested industry personnel interested in the development of best practice guidelines for the consistent achievement of high yielding wheats in a cotton farming system.

A publication titled “Irrigated Wheat- Best Practice Guidelines in Cotton Farming Systems” was generated as part of the project outlining a system that will more

consistently allow producers in the northern growing region of NSW, and Southern Queensland, the ability to achieve high yielding, high quality wheats in an irrigated cotton farming system.



Cotton Catchment Communities CRC

FINAL REPORT

Part 1 - Summary Details

Cotton CRC Project Number: 1.04.11.06

Project Title: Irrigation Scheduling in Wheat

Project Commencement Date: 1/12/2010 **Project Completion Date:** 31/05/2012

Cotton CRC Program: The Farm

Part 2 - Contact Details

Administrator: Justin Harsdorf
Organisation: CSIRO Sustainable Agriculture Flagship
Postal Address: GPO Box 1700, Canberra, ACT, 2601
Ph: (02) 6246 4044 **Fax:** **E-mail:** justin.harsdorf@csiro.au

Principal Researcher: Allan Peake, Project Leader
Organisation: CSIRO Ecosystems Sciences
Postal Address: PO Box 102, Toowoomba, QLD, 4350
Ph: (07) 4688 1137 **Fax:** **E-mail:** allan.peake@csiro.au

Supervisor: Graham Bonnett, Theme Leader
Organisation: CSIRO Sustainable Agriculture Flagship
Postal Address: CSIRO Plant Industry, Queensland Bioscience Precinct, 306 Carmody Road, ST LUCIA QLD 4067

Ph: (07) 3214 2352 **Fax:** **E-mail:** graham.bonnett@csiro.au

Signature of Research Provider Representative: _____



FINAL REPORT

Part 1 - Summary Details

Cotton CRC Project Number: 1.04.11.06

Project Title: Irrigation Scheduling in Wheat

Project Commencement Date: 1/12/2010 **Project Completion Date:** 31/05/2012

Cotton CRC Program: The Farm

Part 2 – Contact Details

Administrator: Justin Harsdorf
Organisation: CSIRO Sustainable Agriculture Flagship
Postal Address: GPO Box 1700, Canberra, ACT, 2601
Ph: (02) 6246 4044 **Fax:** **E-mail:** justin.harsdorf@csiro.au

Principal Researcher: Allan Peake, Project Leader
Organisation: CSIRO Ecosystems Sciences
Postal Address: PO Box 102, Toowoomba, QLD, 4350
Ph: (07) 4688 1137 **Fax:** **E-mail:** allan.peake@csiro.au

Supervisor: Graham Bonnett, Theme Leader
Organisation: CSIRO Sustainable Agriculture Flagship
Postal Address: CSIRO Plant Industry, Queensland Bioscience Precinct, 306 Carmody Road, ST LUCIA QLD 4067

Ph: (07) 3214 2352 **Fax:** **E-mail:** graham.bonnett@csiro.au

Signature of Research Provider Representative: _____

Part 3 – IP and In-kind

Since the November 2011 6 monthly report, please outline the additional IP and in-kind that has been generated in the project.

1. Intellectual Property developed within the project.

(What Know-How (New Ideas), Confidential Information, Copyright, Patents or Provisional Patents, Registered Design, Trade Secrets or Trademarks have come from this project to date)

Not Applicable

2. Project In-kind

(Grower Consultant Ginner or Grower Group In-Kind: Are you conducting part of your project on a cotton farm or in conjunction with an in-kind contribution from a consultant, ginner or Grower Group? Please supply group name - Number of persons involved per week and the number of hours per week involved.)

Not Applicable

Part 4 – Final Report Guide (due at end date of project or 31st May 2012)

(The points below are to be used as a guideline when completing your final report.)

Background

1. Outline the background to the project.

Objectives

2. List the project objectives and the extent to which these have been achieved.

Methods

3. Detail the methodology and justify the methodology used. Include any discoveries in methods that may benefit other related research.

Results

4. Detail and discuss the results for each objective including the statistical analysis of results.

Outcomes

5. Describe how the project's outputs will contribute to the planned outcomes identified in the project application. Describe the planned outcomes achieved to date.
6. Please describe any:-
 - a) technical advances achieved (eg commercially significant developments, patents applied for or granted licenses, etc.);
 - b) other information developed from research (eg discoveries in methodology, equipment design, etc.); and
 - c) required changes to the Intellectual Property register.

Conclusion

7. Provide an assessment of the likely impact of the results and conclusions of the research project for the cotton industry. What are the take home messages?

Extension Opportunities

8. Detail a plan for the activities or other steps that may be taken:
 - (a) to further develop or to exploit the project technology.
 - (b) for the future presentation and dissemination of the project outcomes.
 - (c) for future research.

Publications

A. Publications relevant to this project.

Peer reviewed articles / books

Non-peered reviewed articles

Presentations (conference, field days, workshops etc)

B. All other publications by project team during this period.

Peer reviewed articles / books

Non-peered reviewed articles

(NB: Where possible, please provide a copy of any publication/s)

C. Have you developed any online resources and what is the website address?

Part 5 – Final Report Executive Summary

Provide a one page Summary of your research that is not commercial in confidence, and that can be published on the World Wide Web. Explain the main outcomes of the research and provide contact details for more information. It is important that the Executive Summary highlights concisely the key outputs from the project and, when they are adopted, what this will mean to the cotton industry.

Irrigation Scheduling in Wheat

A final report prepared for the
Cotton Catchment Communities CRC.

Allan S Peake

Citation

Peake AS (2012). Irrigation Scheduling in Wheat. A final report prepared for the Cotton Catchment Communities CRC. CSIRO Sustainable Agriculture Flagship, Canberra, ACT.

Copyright and disclaimer

© 2012 CSIRO To the extent permitted by law, all rights are reserved and no part of this publication covered by copyright may be reproduced or copied in any form or by any means except with the written permission of CSIRO.

Important disclaimer

CSIRO advises that the information contained in this publication comprises general statements based on scientific research. The reader is advised and needs to be aware that such information may be incomplete or unable to be used in any specific situation. No reliance or actions must therefore be made on that information without seeking prior expert professional, scientific and technical advice. To the extent permitted by law, CSIRO (including its employees and consultants) excludes all liability to any person for any consequences, including but not limited to all losses, damages, costs, expenses and any other compensation, arising directly or indirectly from using this publication (in part or in whole) and any information or material contained in it.

Acknowledgements

This study was funded by the Cotton Catchment Communities CRC.

Executive summary

On irrigation farms in northern New South Wales and Queensland, wheat has traditionally been used as a beneficial dryland rotation crop that improves the yield and profitability of subsequent cotton crops. However recent grain price fluctuations have led irrigated enterprises to grow high yielding, irrigated grain crops as a tactical response to water availability and favourable commodity prices. Therefore there is a need for improved advice to irrigated growers who have traditionally grown cotton, and have little experience in producing high yielding, irrigated grain crops. This project aimed to assist irrigated growers in the northern region by producing irrigation scheduling recommendations for irrigated wheat, in conjunction with the CCC CRC 'High Yield Grains in Cotton Farming Systems' (HYGICFS) project.

It was decided to use the APSIM (Agricultural Production Systems Simulator) model to generate irrigation scheduling recommendations for irrigated wheat growing in the northern region, as it is a more cost effective method than wide-scale, multi-year field experiments. However, this first meant that APSIM needed to be 'validated' – that is, the model had to be shown to accurately simulate irrigated wheat production.

Field trials were conducted in 2010 and 2011, to compare the effect of different irrigation strategies on wheat yield. APSIM adequately predicted crop yield and water-use, and it was therefore appropriate to use APSIM to investigate optimum irrigation strategies for irrigated wheat farming systems in the northern grains region.

APSIM simulations showed that at drier environments such as Emerald and Goondiwindi, maximum crop water use efficiency (CWUE) was attained on crops that received two in-crop irrigations, when there were high levels of soil moisture available at sowing (which may require an additional irrigation at sowing). At the wetter environment (Gunnedah), maximum CWUE was achieved with one in-crop irrigation, also assuming high levels of soil moisture available at sowing. Having less soil water at sowing (or low in-crop rainfall) will increase the amount of in-crop irrigation required to achieve maximum CWUE.

Growers wanting to maximise return on a single, 1 ML/ha irrigation, are recommended to apply that irrigation during the period between early stem-elongation through to flag leaf emergence. However, the timing of in-crop rainfall, and the amount of soil water at sowing will affect the optimum timing of this irrigation from season to season.

Through these improved irrigation scheduling guidelines for wheat, irrigated farming systems now have greater capacity to adapt to a variable climate, by having improved ability to grow wheat when the timing of water availability and commodity pricing means it is a profitable option.

Introduction

Wheat is nearly exclusively a dryland crop in the northern grains region of Australia (bounded in the south by Dubbo in New South Wales, and in the north by Emerald in central Queensland (GRDC 2010)). Cotton has traditionally been the favoured irrigation crop in the region, due to its higher profitability in comparison to wheat. In seasons of favourable water supply and prices, up to 400,000 ha of cotton has been produced in these irrigated areas (ANRA 2009). Wheat is commonly considered a beneficial dryland rotation crop that improves the yield and profitability of subsequent cotton crops (Hulugalle *et al.* 2006), through improved soil structure and decreased inoculum of cotton-specific diseases in the soil (Hulugalle *et al.* 1999 & 2006).

In 2007/ 2008, grain prices improved to record levels in response to socioeconomic influences in Asia, an international push for bio-energy crops, and the impact of local droughts. With wheat prices nearly 300% of those received just three years earlier (Brown *et al.* 2008) high levels of optimism were generated for future grain prices. This increase in agricultural commodity prices sparked increased investment in grain farming from the corporate sector, and initiated a paradigm shift amongst the irrigated farming community of the northern grains region. These irrigated enterprises no longer viewed themselves as cotton growers, but rather, commodity growers needing to make the greatest return on the available irrigation water. And while grain prices have since eased, demand for food grain is predicted to increase over the next 20-50 years as the world's population expands (FAO 2006; UN 2007). Demand-driven increases in grain prices are therefore likely to result in increased irrigated wheat production across the northern region.

Therefore there is a need for improved advice to irrigated crop producers that have traditionally grown cotton, but are now commodity producers who need to make tactical responses to water availability and commodity prices in order to maximise on-farm water use efficiency. This project aimed to assist irrigated growers in the northern region by producing irrigation scheduling recommendations for irrigated wheat, in conjunction with the CCC CRC 'High Yield Grains in Cotton Farming Systems' (HYGICFS) project.

Specific project objectives

The primary objectives of the research project were as follows:

- (1) The farming systems simulator APSIM (Agricultural Production Systems Simulator) will be validated using on-farm data to ensure it can appropriately simulate irrigated wheat production.
- (2) APSIM will be used to develop recommendations on irrigation scheduling for wheat, as well as the best way to allocate water to irrigated wheat production in water-limiting scenarios.
- (3) Recommendations will be incorporated into the 'Best Practice Guide' to irrigated wheat production, prepared by the Achievable Yields for Irrigated Grains project and the HYGICFS project team.

Summary of achievements against project objectives

(1) APSIM Validated for irrigated wheat growing

The APSIM model was successfully validated on wheat yield and water use data sets developed in conjunction with the HYGICFS project, from furrow irrigated experiments at ACRI in 2010 and 2011. APSIM successfully simulated yield in both years, and successfully simulated water use in 2011. Water use data from 2010 was not used, as the growing season rainfall was very high, and little discrimination was observed between irrigation treatments.

(2) APSIM used to develop irrigation scheduling recommendations, including the best way to allocate water in water-limiting scenarios.

Irrigation scheduling recommendations were developed using APSIM for two scenarios: (1) maximising crop water use efficiency, and (2) maximising yield from a single 1.1 ML irrigation. The second scenario was developed to cater for growers wishing to maximise the area sown to wheat, to allow them to maximise the area benefiting from a cotton rotation crop. Maximum crop water use efficiency was achieved in simulation experiments where one or two in-crop irrigations of 0.9 ML/ha were applied, and the field had high levels of sowing soil moisture (whether from pre-sowing rainfall or an irrigation at or near sowing). Average maximum yields in response to a single 1.1 ML/ha irrigation were attained when the irrigation was applied during stem elongation, or flag leaf emergence.

(3) Recommendations incorporated into the 'Best Practice Guide for Irrigated Wheat Production' prepared by the HYGICFS project.

The irrigation scheduling recommendations were included in the best practice guide. An additional achievement was the provision of additional graphs to demonstrate other points made by HYGICFS team members in the irrigation scheduling chapter.

Methodology

Overview

Developing irrigation scheduling techniques using field experimentation alone is an expensive, long term process. Variability between seasons and soil types ensures that experiments must be conducted across a large number of sites and seasons in order to sample the full range of conditions that can be experienced by a wheat crop in the northern grains region.

Researchers around the world now realise that simulation modelling has benefits over the traditional field experimentation approach to developing irrigation scheduling recommendations. Lobell and Ortiz-Monasterio (2006) found that the modelling approach was beneficial for optimising system water use efficiency, and allowed the user to examine irrigation management regimes for multiple environments, soil types and climatic conditions. In a review of irrigation management techniques in water scarce environments, Pereira *et al.* (2002) stated: *“More research approaches are required to relate yield responses with gross margin or revenue responses to water deficits. The development of decision support tools integrating irrigation simulation models, namely for extrapolating field trials data, economic evaluation and decision tools should be useful to base the appropriate irrigation management decisions for water scarcity conditions”*. Additionally, the crop modelling approach can be used to demonstrate the level of risk associated with different strategies, by using historical weather data to generate probability distributions of production and profitability (e.g. Hammer *et al.* 1996; Hochman *et al.* 2009; Philp *et al.* 2010).

It was therefore decided to use the APSIM model to generate irrigation scheduling recommendations for irrigated wheat growing in the northern region. A key component of simulation model experiments, however, is that the model must first be ‘validated’ – that is, the model needs to accurately simulate the system being investigated.

Field Experiments

Field experiments were designed in conjunction with staff from NSW DPI, and conducted as part of the HYGICFS project. The field experiments consisted of irrigation scheduling experiments, where wheat plots experienced different levels of irrigation during the season. Experiments were conducted at the Australian Cotton Research Institute (ACRI) at Narrabri in 2010 and 2011. Three irrigation treatments were applied in 2010, and five irrigation treatments were applied in 2011. Irrigation treatments ranged from dryland to fully irrigated, and were conducted in furrow irrigated fields, with the wheat sown on 2m beds across three replicates.

Soil moisture was monitored using a neutron moisture meter (NMM) in each year, except for the surface layer (0-15cm) which was monitored using a Theta probe. Calibration curves for both the NMM and Theta probe were obtained using cores taken in wet and dry parts of the paddock.

Soil characterisation data (Drained Upper Limit or DUL, and Crop Lower Limit or CLL) were obtained as described in Dalgliesh and Foale (1998) for DUL, and by using the lowest soil moisture readings obtained from the NMM, for CLL.

APSIM Validation

Experimental results from the field trials were compared to APSIM simulations of each experimental treatment. Measured soil water and nitrogen at sowing, sowing dates, plant populations, meteorological data, irrigation dates and volumes for each field treatment were used to parameterise each simulation as appropriate. The APSIM ‘skiprow’ logic in the sowing manager was set at 0.3, to simulate the decreased light interception by the irrigation beds.

APSIM Irrigation Scheduling Experiments

APSIM was then used to conduct long term studies on different irrigation strategies using 40 years of weather data from the SILO database (Jeffrey *et al.* 2001). Irrigation scheduling recommendations were developed using APSIM for two scenarios: (1) maximising crop water use efficiency, and (2) maximising yield from a single 1 ML irrigation. The second scenario was developed to cater for growers wishing to increase the area sown to wheat, to allow them to maximise the area benefiting from a cotton rotation crop. Simulations assumed that yields are not limited by pests, diseases, frosts, nutrition or lodging. For agronomic advice to avoid lodging in a high yielding wheat crop, growers are referred to Peake *et al.* (2012).

Strategies to achieve maximum crop water use efficiency (CWUE)

CWUE is defined as the amount of grain produced per unit of water used by the crop or lost to the farm through drainage and evaporation. A series of APSIM simulations were conducted at three locations (Goondiwindi, Gunnedah and Emerald) which investigated CWUE when one, two, three, or four irrigations were applied during the season. The first irrigation occurred at sowing to raise the soil moisture to 2/3 of Plant Available Water Capacity. The remaining irrigations were 0.9 ML/ha irrigation events that occurred when the soil water deficit of the top 120cm of soil was greater than 0.9 ML/ha. The in-crop irrigations were applied at 80% application efficiency, such that 20% of each irrigation was assumed to be lost due to drainage. Representative APSIM soil types (Peake *et al.* 2010) were used for each location, with Gunnedah simulations using Typical Vertosol #3 (PAWC = 255mm), while the Emerald and Goondiwindi simulations used Typical Vertosol #7 (PAWC = 204mm). The cultivar used in these simulations was Kennedy.

Maximising yield from a single irrigation.

A series of APSIM simulations were conducted at three locations (Goondiwindi, Gunnedah and Emerald) which investigated how the timing of one in-crop irrigation affected grain yield. Simulations applied a single, 1.1 ML/ha irrigation at specific growth stages if the soil water deficit to 120cm was greater than 1.1 ML/ha. If the soil water deficit was not large enough at that specific growth stage, the irrigation was delayed until the soil water deficit was equal to (or greater than) 1.1 ML/ha. The soil types and varieties used for these simulations were the same as those used in the CWUE simulations described previously.

Results

APSIM Validation

The APSIM validation demonstrated that APSIM could accurately simulate furrow irrigated wheat production for both yield and water use. Figure 1 shows simulated vs. observed grain yield from different irrigation treatments in 2010 and 2011, where linear regression explained 71% of the variation. These results were achieved without significant incidences of lodging, due in part to specific canopy management techniques that were employed to reduce lodging risk (as described in Peake *et al.* 2012). Heavily lodged crops experience yield reductions that are unable to be predicted by APSIM (Peake, 2009) therefore it is important to note that paddocks must be managed to avoid lodging (as well as disease, pest and frost damage) in order to be able to achieve the yields predicted by APSIM.

Figure 2 demonstrates that APSIM successfully simulated crop water use in 2011. Additional data obtained from monitoring of 4 commercial paddocks in 2008 also showed that APSIM adequately predicted crop water use (Peake, 2009). It is therefore appropriate to use APSIM to estimate both yield and water use for irrigated wheat farming systems in the northern grains region.




FIGURE 1: Simulated vs observed Grain Yield for a range of irrigation treatments at ACRI, 2010 and 2011.

APSIM Irrigation Scheduling Experiments

The APSIM simulation experiments found that on average across the 40 year period studied, maximum CWUE under irrigation was achieved when good soil moisture is available at sowing (or an irrigation had been applied at sowing) and then one or two supplementary spring irrigations were applied. The optimum number of irrigations was one irrigation at Gunnedah and two irrigations at Emerald and Goondiwindi (Figure 3). These results are related to the annual rainfall at these locations, with an additional irrigation required to maximise CWUE at the drier locations (Emerald and Goondiwindi).




FIGURE 2: APSIM simulated Plant Available Water (PAW – solid line) and measured PAW (red squares) for a range of irrigation wheat treatments at ACRI 2011 (a) irrigation at sowing (b) irrigation at sowing + 1 in-crop irrigation (c) irrigation at sowing + 2 in-crop irrigations (d)) irrigation at sowing + 4 in-crop irrigations (e) sown into stored soil-water at sowing + 4 in-crop irrigations.




Figure 3: APSIM-simulated crop water use efficiency of various irrigation scenarios (using 0.9ML/ha irrigation events at 80% application efficiency) for Emerald, Goondiwindi and Gunnedah.

However, consultants and growers should consider that additional irrigations could still return more than 1 tonne of grain per ML of irrigation water use once the maximum CWUE is achieved (Figure 3). Therefore, growers may achieve maximum profitability by applying additional spring-irrigations. The key consideration for growers is whether the yield gain from extra irrigation of the irrigated cropping area is more valuable than the cost of preparing, sowing and managing the additional paddocks, on which they could use the water as an alternative.

The APSIM simulations also found that the best timing for a single in-crop irrigation of around 1 ML/ha is from early stem-elongation, through to flag leaf emergence (Figure 4). While head emergence is the most sensitive growth stage to a short, severe water stress, the APSIM simulations suggest that the best timing for a single irrigation occurs earlier in the life-cycle of the crop, to spread the water across a number of growth stages and avoid a severe stress at any one growth stage. It should be remembered that these results have been obtained on average, from 40 years of weather data. The best timing of a single irrigation within individual seasons will vary depending on the timing of in-crop rainfall, and stored soil water at sowing.




Figure 4: Average yield after application of a 1 ML/ha irrigation at different growth stages, from a 40 year APSIM irrigated wheat simulation at Goondiwindi, Gunnedah and Emerald.

Conclusions & Outcomes

The APSIM farming systems model was shown to accurately predict wheat yield and water use, when the crops were managed to avoid lodging. The APSIM model was used successfully to generate irrigation scheduling recommendations for irrigated wheat growers across the northern grains region.

For growers wanting to maximise crop water use efficiency (CWUE), APSIM simulations showed that at in drier environments such as Emerald and Goondiwindi, maximum CWUE was attained on crops that received two in-crop irrigations, when there were high levels of soil moisture available at sowing (which may require an additional irrigation at or near sowing). At the wetter environment (Gunnedah), maximum CWUE was achieved with just a single in-crop irrigation, also assuming high levels of soil moisture available at sowing. Having less soil water at sowing (or low in-crop rainfall) will increase the amount of in-crop irrigation required to achieve maximum CWUE.

Growers wanting to maximise return on a single, 1 ML/ha irrigation, are recommended to apply that irrigation during the period between early stem-elongation through to flag leaf emergence. However, the timing of in-crop rainfall, and the amount of soil water at sowing will affect the optimum timing of this irrigation from season to season.

The adoption of these irrigation scheduling techniques will (in the long term) help northern region irrigators improve on-farm economic and environmental sustainability through increased profitability and irrigation WUE, and improve the viability of cotton farming systems.

Through these improved irrigation scheduling guidelines for wheat, irrigated farming systems now have greater capacity to adapt to a variable climate, by being able to make tactical decisions to grow wheat (successfully) when the timing of water availability and commodity pricing means it is a profitable option.

Publications

These irrigation scheduling guidelines have been disseminated to growers as part of the following publication, prepared by the CCC-CRC funded 'High Yielding Grains in Cotton Farming Systems' project:

Jackson R, Harris G, Peake AS (2012) Irrigation. *In* Sykes J (Ed), Irrigated Wheat: Best Practice Guidelines in Cotton Farming Systems. Cotton Catchment Communities CRC, Narrabri, NSW

References

- ANRA (2009) Agriculture - Cotton Industry – Australia: Australian National Resource Atlas, Department of the Environment, Water, Heritage and the Arts, <http://www.anra.gov.au/topics/agriculture/cotton/index.html>
- Brown N, Laffan J, Wight M (2008) High food prices, food security and the international trading system, presented to the Informa National Food Pricing Summit, Sydney 19-30 September 2008 by the Assistant Secretary, Trade and Economic Analysis Branch of the Department of Foreign Affairs and Trade (DFAT), Canberra.
- Dalgliesh NP, Foale MA (1998) 'Soil Matters - Monitoring soil water and nutrients in dryland farming', (CSIRO/Agricultural Production Systems Research Unit, Toowoomba, QLD. Technical Manual - ISBN 0 643 06375 7).
- FAO (2006) World Agriculture: Towards 2030/2050. Interim report, FAO, Rome, Italy.
- GRDC (2010) Panel Biographies - Northern Region Characteristics (GRDC website) <http://www.grdc.com.au/director/about/panels/2008.cfm> accessed January 2010.
- Hammer GL, Holzworth DP, Stone R (1996) The value of skill in seasonal climate forecasting to wheat crop management in a region with high climatic variability. *Australian Journal of Agricultural Research* **47**, 717-737.
- Hochman Z, van Rees H, Carberry PS, Hunt JR, McCown RL, Gartmann A, Holzworth D, van Rees S, Dalgliesh NP, Long W, Peake AS, Poulton PL, McClelland T (2009) Re-inventing model-based decision support with Australian dryland farmers. 4. Yield Prophet[®], helps farmers monitor and manage crops in a variable climate. *Crop & Pasture Science* **60**, 2057–2070.
- Hulugalle NR, Entwistle PC, Cooper SF, Nehl DB, Allen SJ, Finlay LA (1999). Sowing wheat or field pea as rotation crops after irrigated cotton in a grey Vertisol. *Australian Journal of Soil Research* **37**, 867-889.
- Hulugalle NR, Weaver TB, Scott, F (2006) Continuous Cotton and Cotton-Wheat Rotation Effects on Soil Properties and Profitability in an Irrigated Vertisol, *Journal of Sustainable Agriculture* **27** (3), 5-24.
- Jeffrey SJ, Carter JO, Moodie KM, Beswick AR (2001) Using spatial interpolation to construct a comprehensive archive of Australian climate data. *Environmental Modelling and Software*, **16**, 309-330.
- UN 2007. World population prospects: The 2006 revision, Highlights. United Nations, New York.
- Lobell DB, Ortiz-Monasterio JI (2006) Evaluating strategies for improved water use in spring wheat with CERES *Agricultural Water Management* **84**, 249-258.
- Peake AS (2009) Monitoring irrigated wheat in the northern region in 2008. Final Report to GRDC project CSA00015. CSIRO Sustainable Ecosystems, ACT.
- Peake AS, Hochman Z, Dalgliesh NP (2010) A rapid method for estimating plant available water capacity on Vertosols. Proceedings of the 15th Australian Society of Agronomy Conference, Lincoln, New Zealand 25-18 November 2010.
- Peake A.S., Rebetzke G., Chapman S., Dreccer F, McIntyre L and Hundt A (2012) Agronomy for high yielding cereal environments: varieties, agronomic strategies and case studies. In GRDC Northern Region Grains Research Updates, Goondiwindi, 6-7 March, 2012. (Publication pending on GRDC website)
- Pereira LS, Oweis T, Zairi A (2002) Irrigation management under water scarcity. *Agricultural Water Management* **57**, 175–206.
- Philp T, Peake AS, McLean G (2010) Improving profitability of irrigated sorghum through partial irrigation of larger areas. Proceedings of the 2010 Australian Summer Grains Conference, Gold Coast, Queensland, 21-24 June 2010.

CONTACT US

t 1300 363 400
+61 3 9545 2176
e enquiries@csiro.au
w www.csiro.au

YOUR CSIRO

Australia is founding its future on science and innovation. Its national science agency, CSIRO, is a powerhouse of ideas, technologies and skills for building prosperity, growth, health and sustainability. It serves governments, industries, business and communities across the nation.

FOR FURTHER INFORMATION

Division/Unit Name

Allan Peake

t +61 7 4688 1137

e allan.peake@csiro.au

w [http://www.csiro.au/Organisation-Structure/Flagships/
Sustainable-AgricultureFlagship/
PeakeAllan_SAF_People.aspx](http://www.csiro.au/Organisation-Structure/Flagships/Sustainable-AgricultureFlagship/PeakeAllan_SAF_People.aspx)